

Prevalence of Insomnia among Post Covid 19 Elderly Population

Annie Jerin Retna Raj, Kumaresan Abathsagayam*, Surya Vishnuram, Prathap Suganthirababu, Vignesh Srinivasan, Vinodhkumar Ramalingam, & Preethi Gokulan

Saveetha College of Physiotherapy, Saveetha Institute of Medical and Technical Sciences,
Chennai, Tamilnadu, India.

***Email:** kumaresan.scpt@saveetha.com

Abstract

Insomnia is defined as a persistent difficulty initiating or maintaining sleep, or by non-restorative sleep that causes daytime distress. Insomnia can lead to mental health issues such as depression and anxiousness, as well as an elevated risk long-term diseases or conditions like high blood pressure and heart disease. COVID -19 is a significant psychological stressor for patients. Patients are vulnerable to a variety of mental health issues even after they have recovered. The aim and objective of the study is to find out the prevalence of insomnia among post COVID- 19 elderly population. This study is a survey. The study was conducted among 300 participants of post COVID 19 Elderly population. Based upon the inclusion and exclusion criteria participants were selected. The detailed procedure was explained to the participants and a written informed consent form was obtained. The insomnia severity Index scale was circulated among the participants and the obtained data was analyzed. The values were statistically analyzed. This statistical analysis shows that in 1st wave 97 males and 73 females were affected and in 2nd wave 49 males and 71 females were affected. This article concluded that there is prevalence of insomnia among post COVID 19 elderly population.

Keywords

COVID 19, Insomnia, elderly population

Introduction

COVID-19 is a novel coronavirus disease that was initially found in Wuhan, Hubei Province, in December 2019 and has since spread to other parts of China and then to many other nations (Ahn et al., 2019; Kannan et al., 2019). On March 11, 2020, the World Health Organization (WHO) proclaimed COVID-19 a pandemic. More than 15 million people have been infected with COVID-19 as of July 25, 2020, according to the WHO, and 635,173 people had died worldwide. COVID-19 has not only sparked international concern but has also sparked local concern (Xiang et al., 2020). People were also subjected to a tremendous deal of psychological stress because of depression, anxiety, and psychotic symptoms were all recorded during the SARS outbreak, according to previous investigations (Xu et al., 2022; Torales et al., 2022; Huang et al., 2020; Shi

Submission: 2 June 2023; **Acceptance:** 3 July 2023



Copyright: © 2023. All the authors listed in this paper. The distribution, reproduction, and any other usage of the content of this paper is permitted, with credit given to all the author(s) and copyright owner(s) in accordance to common academic practice. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license, as stated in the website: <https://creativecommons.org/licenses/by/4.0/>

et al., 2020; Shi et al., 2020; Maunder et al., 2003; Nickell et al., 2004; Wu et al., 2009). Among older persons and those with comorbidities like as hypertension, cardiovascular illness, diabetes, chronic lung disease, and chronic kidney disease, studies have indicated that this virus causes worse results and a greater mortality rate (Liu et al., 2021) Further-more, few COVID-19 patients died because of the infection exacerbating underlying comorbid health issues, rather than the illness itself.

In Italy and China, data stratification by age revealed a nearly identical case-fatality rate for adults aged 69. Individuals over the age of 70, particularly those over the age of 80, had a higher prevalence in Italy (Xu et al., 2020). In a study of 5700 COVID-19 patients in New York, the in-hospital death rate for adults aged 60–69, 70–79, 80–89, and >90 years was 15.8, 32.2, 54.3, and 52.3 percent, respectively (Mez et al., 2017). These disorders affect a large number of elderly Americans, putting them at a higher risk of infection. Infection with the severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) is linked to an elevated risk of long-term cognitive loss in the elderly. COVID-19 patients, especially those with severe cases, should be closely evaluated for cognitive loss after infection (Xu et al., 2020). COVID-19 was not only an international concern (Xiang et al., 2020) but also has caused a variety of mental health difficulties, particularly among confirmed patients, due to its quick onset, rapid transmission, high mortality, lack of effective therapies, and large-scale isolation measures. During their time in the hospital, they experienced both physical and psychological hardship (Wang et al., 2022).

In a Chinese pilot study, nearly half of the participants rated COVID-19's mental health impact as moderate or severe, and one-third felt anxiety symptoms (Shi et al., 2020). In this issue of Sleep Medicine, Lin and colleagues report on a timely study that examined the early effects of the COVID-19 outbreak on sleep and psychiatric conditions in a huge sample of adults (N = 5641) in China. The main results indicated greater incidence of clinically severe insomnia, as expected (20%) (Morin et al., 2021). We discovered that COVID-19 recovered patients had a low rate of depression and a high rate of insomnia based on the findings. We should pay more attention to their sleeping habits than to their mood (Wang et al., 2022).

Older age, female gender, and co morbid medical and psychiatric conditions are all associated with a higher prevalence of chronic insomnia. Around 40% of persons with insomnia also have a diagnosable psychiatric condition, the most common of which is depression (Roth et al., 2007). In the absence of a clear physical, mental, or substance-related reason, primary insomnia is described as difficulty falling asleep, sustaining sleep, or getting restful sleep, as well as markedly reduced daytime functioning. 40% of insomnia patients are thought to have a psychological disorder.

Depression is the most frequent of these psychiatric diseases, and insomnia is a symptom of both. Estimated prevalence of insomnia in the United States varies from 3.9% to 27.3% and increases with age, reaching 30% to 48% in older adults (McElroy et al., 2021). Adult patients older than the age of 65.6 accounted for more than 80% of all deaths due to COVID 19 (Shahid et al., 2020). Therefore, they are more prone to insomnia.

Methodology

This prevalence study was conducted at Saveetha Medical Hospital. The survey was conducted among the post-COVID-19 elderly population. This was carried out using an insomnia severity index questionnaire. The study was started in February 2022. The demographic details like Name and age gender were collected. People above 60 years of age were selected from both genders. The selected person tested positive for COVID-19 and completed 2 weeks of the recovery period. Participants with cognitive decline, uncooperative people, and elderly people with critical psychological problems were excluded. The detailed procedure has been explained and informed consent is obtained from the participants. All the participants were asked to fill out the insomnia severity index questionnaire. In the first wave, 97 males and 73 females were present and in the second wave 49 males and 71 females were present. The values are tabulated and statistically analyzed.

Table 1. Participants in 1st and 2nd wave

Wave	Gender	Total Number
1 ST Wave	Male	97
	Female	73
2 nd Wave	Male	49
	Female	71

Table 2. Tabulated mean values of insomnia classification

Waves	Gender	No Clinically Significant Insomnia	Sub Threshold Insomnia	Clinical Insomnia (Moderate)	Clinical Insomnia (Severe)
1 st wave	Male	4.285	10.26	17.93	24.933
	Female	2.375	11	17.97	25.58
2 nd wave	Male	3.5	10.77	18.8	23.85
	Female	3.5	10.76	18.84	24.2

Table 3. Tabulated percentage values for insomnia classification

Waves	Gender	No Clinically Significant Insomnia	Sub Threshold Insomnia	Clinical Insomnia (Moderate)	Clinical Insomnia (Severe)
1 st Wave	Male	14.43%	30.92%	31.95%	22.68%
	Female	10.95%	9.5%	58.90%	16.43%
2 nd Wave	Male	16.32%	18.36%	51.02%	14.28%
	Female	8.45%	29.57%	26.76%	35.21%

Results and Discussion

The values were statistically analyzed. This statistical analysis showed that in the 1st wave 97 males and 73 females were affected and in the 2nd wave 49 males and 71 females were affected. The

mean value for males in the first wave for no clinically significant insomnia was 4.285, subthreshold insomnia was 10.26, clinical insomnia (moderate) was 17.93 and clinical insomnia (severe) was 24 and for females for no clinically significant insomnia was 2.375, subthreshold insomnia was 11, clinical insomnia (moderate) was 17.97 and clinical insomnia (severe) was 25.58. The mean value for males in the second wave for no clinically significant insomnia was 3.5, subthreshold insomnia was 10.77, clinical insomnia (moderate) was 18.8 and clinical insomnia (severe) was 23.85 and for females for no clinically significant insomnia was 3.5, subthreshold insomnia was 10.76, clinical insomnia (moderate) was 18.84 and clinical insomnia (severe) was 24.2.

COVID-19 patients had a significant prevalence of insomnia (26.45 percent) two weeks after discharge, according to the findings of the on-site assessment, and had risk factors for developing insomnia (Xu et al., 2022). Many other similar research, have found that individuals with pre-existing disorders and COVID-19 infection have the highest rates of depression and anxiety (Luo et al., 2020; Zhang et al., 2020; Wu et al., 2005). A comparable study found that 18% of 131 individuals experienced depressive symptoms one month following discharge (Lai et al., 2019). A recent mental health assessment of 1257 health care employees revealed a significant prevalence of mental health symptoms (above 70%), with sleeplessness accounting for 34% and depression accounting for 50.4 percent (Huang et al., 2020).

On the other hand, one in every five individuals suffered insomnia, implying that the public's psychological stress may be increased as the epidemic progresses (Tang et al., 2020; Xiang et al., 2020). As a result, the older patients were severely impacted both physically and mentally following the COVID-19 pandemic. According to recent research, COVID-19-infected people had a greater frequency of insomnia and depression as a result of social isolation, uncertainty, and health state (Yang et al., 2020). The perceived danger and worry of spreading the illness to others could be the source of insomnia and sadness. Furthermore, COVID-19's quick transmission, high mortality, and negative news may raise the likelihood of insomnia and depression worsen physical and mental impairment, and further damage patients' daily functions and cognitive capacity (Morin et al., 1993). Fei et al (2021) stated that COVID-19 infected patients after 2 weeks discharge showed 26.45% insomnia. The result of our study concluded that the percentage of males in 1st wave among post COVID-19 elderly population is 54.63% and 75.33% in females. The percentage of males in 2nd wave among post COVID-19 elderly population is 65.3% and 61.97% in females.

Conclusion

The study concluded that males were affected more than females in 1st wave and females were affected more than males in second wave. Insomnia severity index scale was used which showed the results that the prevalence of insomnia is greater in post COVID 19 elderly population.

Acknowledgements

Not applicable.

References

- Ahn, D. G., Shin, H. J., Kim, M. H., Lee, S., Kim, H. S., Myoung, J., ... & Kim, S. J. (2020). Current status of epidemiology, diagnosis, therapeutics, and vaccines for novel coronavirus disease 2019 (COVID-19).
- Bastien, C. H., Vallières, A., & Morin, C. M. (2001). Validation of the Insomnia Severity Index as an outcome measure for insomnia research. *Sleep Medicine*, 2(4), 297–307.
- Deng, J., Zhou, F., Hou, W., Silver, Z., Wong, C. Y., Chang, O., ... & Zuo, Q. K. (2021). The prevalence of depression, anxiety, and sleep disturbances in COVID-19 patients: A meta-analysis. *Annals of the New York Academy of Sciences*, 1486(1), 90–111.
- Huang, Y., & Zhao, N. (2020). Generalized anxiety disorder, depressive symptoms and sleep quality during COVID-19 outbreak in China: A web-based cross-sectional survey. *Psychiatry Research*, 288, 112954.
- Kannan, S. P. A. S., Ali, P. S. S., Sheeza, A., & Hemalatha, K. (2020). COVID-19 (Novel Coronavirus 2019) – Recent trends. *European Review for Medical and Pharmacological Sciences*, 24(4), 2006–2011.
- Lai, J., Ma, S., Wang, Y., Cai, Z., Hu, J., Wei, N., ... & Hu, S. (2020). Factors associated with mental health outcomes among health care workers exposed to coronavirus disease 2019. *JAMA Network Open*, 3(3), e203976.
- Liu, Y. H., Wang, Y. R., Wang, Q. H., Chen, Y., Chen, X., Li, Y., ... & Wang, Y. J. (2021). Post-infection cognitive impairments in a cohort of elderly patients with COVID-19. *Molecular Neurodegeneration*, 16(1), 1–10.
- Luo, M., Guo, L., Yu, M., Jiang, W., & Wang, H. (2020). The psychological and mental impact of coronavirus disease 2019 (COVID-19) on medical staff and general public: A systematic review and meta-analysis. *Psychiatry Research*, 291, 113190.
- Maunder, R., Hunter, J., Vincent, L., Bennett, J., Peladeau, N., Leszcz, M., ... & Mazzulli, T. (2003). The immediate psychological and occupational impact of the 2003 SARS outbreak in a teaching hospital. *CMAJ*, 168(10), 1245–1251.
- Mendes, A., Serratrice, C., Herrmann, F. R., Genton, L., Périvier, S., Scheffler, M., ... & Zekry, D. (2020). Predictors of in-hospital mortality in older patients with COVID-19: The COVIDAge study. *Journal of the American Medical Directors Association*, 21(11), 1546–1554.
- Mez, J., Daneshvar, D. H., Kiernan, P. T., Abdolmohammadi, B., Alvarez, V. E., Huber, B. R., ... & McKee, A. C. (2017). Clinicopathological evaluation of chronic traumatic encephalopathy in players of American football. *JAMA*, 318(4), 360–370.
- Morin, C. M. (1993). *Insomnia: Psychological assessment and management*. Guilford Press.
- Morin, C. M., & Carrier, J. (2021). The acute effects of the COVID-19 pandemic on insomnia and psychological symptoms. *Sleep Medicine*, 77, 346.
- Nickell, L. A., Crighton, E. J., Tracy, C. S., Al-Enazy, H., Bolaji, Y., Hanjrah, S., ... & Upshur, R. E. (2004). Psychosocial effects of SARS on hospital staff: Survey of a large tertiary care institution. *CMAJ*, 170(5), 793–798.
- Patel, D., Steinberg, J., & Patel, P. (2018). Insomnia in the elderly: A review. *Journal of Clinical Sleep Medicine*, 14(6), 1017–1024.
- Roth, T. (2007). Insomnia: Definition, prevalence, etiology, and consequences. *Journal of Clinical Sleep Medicine*, 3(5 Suppl), S7–S10.

- Shahid, Z., Kalayanamitra, R., McClafferty, B., Kepko, D., Ramgobin, D., Patel, R., ... & Jain, R. (2020). COVID-19 and older adults: What we know. *Journal of the American Geriatrics Society*, 68(5), 926–929.
- Shi, L., Lu, Z. A., Que, J. Y., Huang, X. L., Liu, L., Ran, M. S., ... & Lu, L. (2020). Prevalence of and risk factors associated with mental health symptoms among the general population in China during the coronavirus disease 2019 pandemic. *JAMA Network Open*, 3(7), e2014053.
- Shi, Y., Wang, J., Yang, Y., Wang, Z., Wang, G., Hashimoto, K., ... & Liu, H. (2020). Knowledge and attitudes of medical staff in Chinese psychiatric hospitals regarding COVID-19. *Brain, Behavior, & Immunity–Health*, 4, 100064.
- Tang, W., Hu, T., Hu, B., Jin, C., Wang, G., Xie, C., ... & Xu, J. (2020). Prevalence and correlates of PTSD and depressive symptoms one month after the outbreak of the COVID-19 epidemic in a sample of home-quarantined Chinese university students. *Journal of Affective Disorders*, 274, 1–7.
- Torales, J., O’Higgins, M., Castaldelli-Maia, J. M., & Ventriglio, A. (2020). The outbreak of COVID-19 coronavirus and its impact on global mental health. *International Journal of Social Psychiatry*, 66(4), 317–320.
- Wu, K. K., Chan, S. K., & Ma, T. M. (2005). Posttraumatic stress after SARS. *Emerging Infectious Diseases*, 11(8), 1297.
- Wu, P., Fang, Y., Guan, Z., Fan, B., Kong, J., Yao, Z., ... & Hoven, C. W. (2009). The psychological impact of the SARS epidemic on hospital employees in China: Exposure, risk perception, and altruistic acceptance of risk. *The Canadian Journal of Psychiatry*, 54(5), 302–311.
- Xiang, Y. T., Li, W., Zhang, Q., Jin, Y., Rao, W. W., Zeng, L. N., ... & Hall, B. J. (2020). Timely research papers about COVID-19 in China. *The Lancet*, 395(10225), 684–685.
- Xiang, Y. T., Yang, Y., Li, W., Zhang, L., Zhang, Q., Cheung, T., & Ng, C. H. (2020). Timely mental health care for the 2019 novel coronavirus outbreak is urgently needed. *The Lancet Psychiatry*, 7(3), 228–229.
- Xu, F., Wang, X., Yang, Y., Zhang, K., Shi, Y., Xia, L., ... & Liu, H. (2022). Depression and insomnia in COVID-19 survivors: A cross-sectional survey from Chinese rehabilitation centers in Anhui province. *Sleep Medicine*, 91, 161–165.
- Yang, Y., Li, W., Zhang, Q., Zhang, L., Cheung, T., & Xiang, Y. T. (2020). Mental health services for older adults in China during the COVID-19 outbreak. *The Lancet Psychiatry*, 7(4), e19.
- Zhang, J., Lu, H., Zeng, H., Zhang, S., Du, Q., Jiang, T., & Du, B. (2020). The differential psychological distress of populations affected by the COVID-19 pandemic. *Brain, Behavior, and Immunity*, 87, 49.